

A CLASS OF HARMONIC FUNCTIONS WITH REAL COEFFICIENTS DEFINED BY CONVOLUTION

SUMIT NAGPAL AND V. RAVICHANDRAN

ABSTRACT. For given two harmonic functions Φ and Ψ with real coefficients in the open unit disk $\mathbb{D}$, we study a class of harmonic functions $f(z) = z - \sum_{n=2}^{\infty} a_n z^n + \sum_{n=1}^{\infty} b_n \bar{z}^n$ ($a_n, b_n \geq 0$) satisfying

$$\operatorname{Re} \frac{(f * \Phi)(z)}{(f * \Psi)(z)} > \alpha \quad (0 \leq \alpha < 1, z \in \mathbb{D});$$

$*$ being the harmonic convolution. Coefficient inequalities, growth and covering theorems, as well as closure theorems are determined. The results obtained extend several known results as special cases. In addition, we study the class of harmonic functions f that satisfy $\operatorname{Re} f(z)/z > \alpha$ ($0 \leq \alpha < 1, z \in \mathbb{D}$).

1. INTRODUCTION

Let $\mathcal{H}$ denote the class of all complex-valued harmonic functions f in the open unit disk $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ normalized by $f(0) = 0 = f_z(0) - 1$. Such functions can be written in the form $f = h + \bar{g}$, where

$$(1.1) \quad h(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad \text{and} \quad g(z) = \sum_{n=1}^{\infty} b_n z^n$$

are analytic in $\mathbb{D}$. We call h the analytic part and g the co-analytic part of f . Let $\mathcal{S}_H$ be the subclass of $\mathcal{H}$ consisting of univalent and sense-preserving functions. In 1984, Clunie and Sheil-Small [4] initiated the study of the class $\mathcal{S}_H$ and its subclasses. Recent works in this direction include those of [15, 16].

For analytic functions $\phi(z) = z + \sum_{n=2}^{\infty} a_n z^n$ and $\psi(z) = z + \sum_{n=2}^{\infty} A_n z^n$, their convolution (or Hadamard product) is defined as $(\phi * \psi)(z) = z + \sum_{n=2}^{\infty} a_n A_n z^n$, $z \in \mathbb{D}$. In the harmonic case, with $f = h + \bar{g}$ and $F = H + \bar{G}$, their harmonic convolution is defined as $f * F = h * H + \overline{g * G}$.

Let $\mathcal{TH}$ be the subclass of $\mathcal{H}$ consisting of functions $f = h + \bar{g}$ so that h and g take the form

$$(1.2) \quad h(z) = z - \sum_{n=2}^{\infty} a_n z^n, \quad g(z) = \sum_{n=1}^{\infty} b_n z^n \quad (a_n, b_n \geq 0).$$

Motivated with the work of Juneja *et al.* [11] (in the analytic case) and making use of the convolution structure for harmonic mappings, we study a new subclass $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ introduced in the following:

2010 *Mathematics Subject Classification.* 30C45, 30C50.

Key words and phrases. harmonic mappings; convolution; convex and starlike functions.

The research work of the first author is supported by research fellowship from Council of Scientific and Industrial Research (CSIR), New Delhi.

Definition 1.1. Suppose that $i, j \in \{0, 1\}$. Let the functions Φ_i, Ψ_j given by

$$\Phi_i(z) = z + \sum_{n=2}^{\infty} p_n z^n + (-1)^i \sum_{n=1}^{\infty} q_n \bar{z}^n \quad \text{and} \quad \Psi_j(z) = z + \sum_{n=2}^{\infty} u_n z^n + (-1)^j \sum_{n=1}^{\infty} v_n \bar{z}^n$$

are harmonic in $\mathbb{D}$ with $p_n \geq u_n \geq 0$ ($n = 2, 3, \dots$) and $q_n \geq v_n \geq 0$ ($n = 1, 2, \dots$). Then a function $f \in \mathcal{H}$ is said to be in the class $\mathcal{H}(\Phi_i, \Psi_j; \alpha)$ if and only if

$$(1.3) \quad \operatorname{Re} \frac{(f * \Phi_i)(z)}{(f * \Psi_j)(z)} > \alpha \quad (z \in \mathbb{D}),$$

where $*$ denotes the harmonic convolution as defined above and $0 \leq \alpha < 1$. Further we define the class $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ by

$$\mathcal{TH}(\Phi_i, \Psi_j; \alpha) = \mathcal{H}(\Phi_i, \Psi_j; \alpha) \cap \mathcal{TH}.$$

The family $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ includes a variety of well-known subclasses of harmonic functions as well as many new ones. For example

$$\mathcal{TH} \left(\frac{z}{(1-z)^2} - \frac{\bar{z}}{(1-\bar{z})^2}, \frac{z}{1-z} + \frac{\bar{z}}{1-\bar{z}}; \alpha \right) \equiv \mathcal{TS}_H^*(\alpha);$$

and

$$\mathcal{TH} \left(\frac{z+z^2}{(1-z)^3} + \frac{\bar{z}+\bar{z}^2}{(1-\bar{z})^3}, \frac{z}{(1-z)^2} - \frac{\bar{z}}{(1-\bar{z})^2}; \alpha \right) \equiv \mathcal{TK}_H(\alpha)$$

are the classes of sense-preserving harmonic univalent functions f which are starlike of order α ($0 \leq \alpha < 1$) and convex of order α ($0 \leq \alpha < 1$) respectively (see [8, 9, 16]). These classes were studied by Silverman and Silvia [18] for the case $\alpha = 0$. Recall that starlike functions of order α and convex functions of order α are characterized by the conditions

$$\frac{\partial}{\partial \theta} \arg f(re^{i\theta}) > \alpha \quad \text{and} \quad \frac{\partial}{\partial \theta} \left(\arg \left\{ \frac{\partial}{\partial \theta} f(re^{i\theta}) \right\} \right) > \alpha, \quad (0 \leq \theta < 2\pi, 0 < r < 1)$$

respectively. In the similar fashion, it is easy to see that the subclasses of $\mathcal{TH}$ introduced by Ahuja *et al.* [1, 2], Dixit *et al.* [5, 6], Frasin [7], Joshi [10], Murugusundaramoorthy *et al.* [12, 13, 14] and Yalçın *et al.* [19, 20] are special cases of our class $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ for suitable choice of the functions Φ_i and Ψ_j .

In Section 2, we obtain the coefficient inequalities, growth and covering theorems, as well as closure theorems for functions in the class $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$. Additionally, we present results for the class of harmonic functions f that satisfy $\operatorname{Re} f(z)/z > \alpha$ ($0 \leq \alpha < 1, z \in \mathbb{D}$) in the last section of the paper.

2. THE CLASS $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$

The first theorem of this section provides a sufficient coefficient condition for a function to be in the class $\mathcal{H}(\Phi_i, \Psi_j; \alpha)$.

Theorem 2.1. *Let the function $f = h + \bar{g}$ be such that h and g are given by (1.1). Furthermore, let*

$$(2.1) \quad \sum_{n=2}^{\infty} \frac{p_n - \alpha u_n}{1 - \alpha} |a_n| + \sum_{n=1}^{\infty} \frac{q_n - (-1)^{j-i} \alpha v_n}{1 - \alpha} |b_n| \leq 1$$

where $0 \leq \alpha < 1$, $i, j \in \{0, 1\}$, $p_n \geq u_n \geq 0$ ($n = 2, 3, \dots$) and $q_n \geq v_n \geq 0$ ($n = 1, 2, \dots$). Then $f \in \mathcal{H}(\Phi_i, \Psi_j; \alpha)$.

Proof. Using the fact that $\operatorname{Re} w > \alpha$ if and only if $|w - 1| < |w + 1 - 2\alpha|$, it suffices to show that

$$(2.2) \quad |A(z) + (1 - 2\alpha)B(z)| - |A(z) - B(z)| \geq 0,$$

where

$$A(z) = (f * \Phi_i)(z) = z + \sum_{n=2}^{\infty} a_n p_n z^n + (-1)^i \sum_{n=1}^{\infty} b_n q_n \bar{z}^n$$

and

$$B(z) = (f * \Psi_j)(z) = z + \sum_{n=2}^{\infty} a_n u_n z^n + (-1)^j \sum_{n=1}^{\infty} b_n v_n \bar{z}^n.$$

Substituting for $A(z)$ and $B(z)$ in (2.2) and making use of (2.1) we obtain

$$\begin{aligned} & |A(z) + (1 - 2\alpha)B(z)| - |A(z) - B(z)| \\ &= \left| 2(1 - \alpha)z + \sum_{n=2}^{\infty} (p_n + (1 - 2\alpha)u_n)a_n z^n + (-1)^i \sum_{n=1}^{\infty} (q_n + (-1)^{j-i}(1 - 2\alpha)v_n)b_n \bar{z}^n \right| \\ &\quad - \left| \sum_{n=2}^{\infty} (p_n - u_n)a_n z^n + (-1)^i \sum_{n=1}^{\infty} (q_n - (-1)^{j-i}v_n)b_n \bar{z}^n \right| \\ &\geq 2(1 - \alpha)|z| - \sum_{n=2}^{\infty} (p_n + (1 - 2\alpha)u_n)|a_n||z|^n - \sum_{n=1}^{\infty} (q_n + (-1)^{j-i}(1 - 2\alpha)v_n)|b_n||z|^n \\ &\quad - \sum_{n=2}^{\infty} (p_n - u_n)|a_n||z|^n - \sum_{n=1}^{\infty} (q_n - (-1)^{j-i}v_n)|b_n||z|^n \\ &= 2(1 - \alpha)|z| \left[1 - \sum_{n=2}^{\infty} \frac{p_n - \alpha u_n}{1 - \alpha} |a_n||z|^{n-1} - \sum_{n=1}^{\infty} \frac{q_n - (-1)^{j-i}\alpha v_n}{1 - \alpha} |b_n||z|^{n-1} \right] \\ &> 2(1 - \alpha)|z| \left[1 - \sum_{n=2}^{\infty} \frac{p_n - \alpha u_n}{1 - \alpha} |a_n| - \sum_{n=1}^{\infty} \frac{q_n - (-1)^{j-i}\alpha v_n}{1 - \alpha} |b_n| \right] \geq 0. \end{aligned}$$

The harmonic mappings

$$f(z) = z + \sum_{n=2}^{\infty} \frac{1 - \alpha}{p_n - \alpha u_n} x_n z^n + \sum_{n=1}^{\infty} \frac{1 - \alpha}{q_n - (-1)^{j-i}\alpha v_n} \bar{y}_n \bar{z}^n,$$

where $\sum_{n=2}^{\infty} |x_n| + \sum_{n=1}^{\infty} |y_n| = 1$, show that the coefficient bound given by (2.1) is sharp. $\square$

Remark 2.2. In addition to the hypothesis of Theorem 2.1, if we assume that $p_n \geq n$ ($n = 2, 3, \dots$) and $q_n \geq n$ ($n = 1, 2, \dots$) then it is easy to deduce that $n(1 - \alpha) \leq p_n - \alpha u_n$ ($n = 2, 3, \dots$) and $n(1 - \alpha) \leq q_n - (-1)^{j-i}\alpha v_n$ ($n = 1, 2, \dots$) so that

$$\sum_{n=2}^{\infty} n|a_n| + \sum_{n=1}^{\infty} n|b_n| \leq \sum_{n=2}^{\infty} \frac{p_n - \alpha u_n}{1 - \alpha} |a_n| + \sum_{n=1}^{\infty} \frac{q_n - (-1)^{j-i}\alpha v_n}{1 - \alpha} |b_n| \leq 1.$$

By [9, Theorem 1, p. 472], $f \in \mathcal{S}_H$ and maps $\mathbb{D}$ onto a starlike domain.

Let $F(a, b, c; z)$ be the Gaussian hypergeometric function defined by

$$F(a, b, c; z) := \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n (1)_n} z^n, \quad z \in \mathbb{D}$$

where a, b, c are complex numbers with $c \neq 0, -1, -2, \dots$ and $(\theta)_n$ is the Pochhammer symbol: $(\theta)_0 = 1$ and $(\theta)_n = \theta(\theta+1)\dots(\theta+n-1)$ for $n = 1, 2, \dots$. Theorem 2.1 gives a sufficient condition for the harmonic function $\phi_1 + \bar{\phi}_2$ to be in the class $\mathcal{H}(\Phi_i, \Psi_j; \alpha)$ where $\phi_1(z) \equiv \phi_1(a_1, b_1, c_1; z)$ and $\phi_2(z) \equiv \phi_2(a_2, b_2, c_2; z)$ are the hypergeometric functions defined by

$$(2.3) \quad \phi_1(z) := zF(a_1, b_1, c_1; z) \quad \text{and} \quad \phi_2(z) := F(a_2, b_2, c_2; z) - 1.$$

Corollary 2.3. *Let $a_k, b_k > 0$, $c_k > a_k + b_k + 1$ for $k = 1, 2$. Furthermore, let*

$$\sum_{n=2}^{\infty} \frac{p_n - \alpha u_n}{1 - \alpha} \frac{(a_1)_{n-1} (b_1)_{n-1}}{(c_1)_{n-1} (1)_{n-1}} + \sum_{n=1}^{\infty} \frac{q_n - (-1)^{j-i} \alpha v_n}{1 - \alpha} \frac{(a_2)_n (b_2)_n}{(c_2)_n (1)_n} \leq 1$$

where $0 \leq \alpha < 1$, $i, j \in \{0, 1\}$, $p_n \geq u_n \geq 0$ ($n = 2, 3, \dots$) and $q_n \geq v_n \geq 0$ ($n = 1, 2, \dots$). Then $\phi_1 + \bar{\phi}_2 \in \mathcal{H}(\Phi_i, \Psi_j; \alpha)$, ϕ_1 and ϕ_2 being given by (2.3).

It is worth to remark that Theorems 2.2, 2.4 and 2.11 of [3] are particular cases of Corollary 2.3. We next show that the coefficient condition (2.1) is also necessary for functions in $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$.

Theorem 2.4. *Let the function $f = h + \bar{g}$ be such that h and g are given by (1.2). Then $f \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ if and only if*

$$(2.4) \quad \sum_{n=2}^{\infty} \frac{p_n - \alpha u_n}{1 - \alpha} a_n + \sum_{n=1}^{\infty} \frac{q_n - (-1)^{j-i} \alpha v_n}{1 - \alpha} b_n \leq 1$$

where $0 \leq \alpha < 1$, $i, j \in \{0, 1\}$, $p_n \geq u_n \geq 0$ ($n = 2, 3, \dots$) and $q_n \geq v_n \geq 0$ ($n = 1, 2, \dots$).

Proof. The sufficient part follows by Theorem 2.1 upon noting that $\mathcal{TH}(\Phi_i, \Psi_j; \alpha) \subset \mathcal{H}(\Phi_i, \Psi_j; \alpha)$. For the necessary part, let $f \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$. Then (1.3) yields

$$\begin{aligned} & \operatorname{Re} \left(\frac{(f * \Phi_i)(z)}{(f * \Psi_j)(z)} - \alpha \right) \\ &= \operatorname{Re} \left(\frac{(1 - \alpha)z + \sum_{n=2}^{\infty} (p_n - \alpha u_n) a_n z^n + (-1)^i \sum_{n=1}^{\infty} (q_n - (-1)^{j-i} \alpha v_n) b_n \bar{z}^n}{z + \sum_{n=2}^{\infty} a_n u_n z^n + (-1)^j \sum_{n=1}^{\infty} b_n v_n \bar{z}^n} \right) \\ &\geq \frac{(1 - \alpha) - \sum_{n=2}^{\infty} (p_n - \alpha u_n) a_n |z|^{n-1} - \sum_{n=1}^{\infty} (q_n - (-1)^{j-i} \alpha v_n) b_n |z|^{n-1}}{1 + \sum_{n=2}^{\infty} a_n u_n |z|^{n-1} + \sum_{n=1}^{\infty} b_n v_n |z|^{n-1}} > 0. \end{aligned}$$

The above inequality must hold for all $z \in \mathbb{D}$. In particular, choosing the values of z on the positive real axis and letting $z \rightarrow 1^-$ we obtain the required condition (2.4). $\square$

Theorem 2.4 immediately yields the following two corollaries.

Corollary 2.5. For $f = h + \bar{g} \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ where h and g are given by (1.2), we have

$$a_n \leq \frac{1 - \alpha}{p_n - \alpha u_n} \quad (n = 2, 3, \dots) \quad \text{and} \quad b_n \leq \frac{1 - \alpha}{q_n - (-1)^{j-i} \alpha v_n} \quad (n = 1, 2, \dots);$$

the result being sharp, for each n .

Corollary 2.6. Suppose that $\Phi(z) = z + \sum_{n=2}^{\infty} p_n z^n + \sum_{n=1}^{\infty} q_n \bar{z}^n$ ($p_n, q_n \geq 0$) is a harmonic function in $\mathbb{D}$ and $f = h + \bar{g} \in \mathcal{TH}$ where h and g are given by (1.2). Then

$$\operatorname{Re} \frac{(f * \Phi)(z)}{z} > \alpha \quad (z \in \mathbb{D}) \quad \Leftrightarrow \quad \sum_{n=2}^{\infty} \frac{p_n}{1 - \alpha} a_n + \sum_{n=1}^{\infty} \frac{q_n}{1 - \alpha} b_n \leq 1.$$

By making use of Theorem 2.4, we obtain the following growth estimate for functions in the class $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$.

Theorem 2.7. Let $f \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$, $\sigma_n = p_n - \alpha u_n$ ($n = 2, 3, \dots$) and $\Gamma_n = q_n - (-1)^{j-i} \alpha v_n$ ($n = 1, 2, \dots$). If $\{\sigma_n\}$ and $\{\Gamma_n\}$ are non-decreasing sequences, then

$$|f(z)| \leq (1 + b_1)|z| + \frac{1 - \alpha}{\eta} \left(1 - \frac{q_1 - (-1)^{j-i} \alpha v_1}{1 - \alpha} b_1 \right) |z|^2,$$

and

$$|f(z)| \geq (1 - b_1)|z| - \frac{1 - \alpha}{\eta} \left(1 - \frac{q_1 - (-1)^{j-i} \alpha v_1}{1 - \alpha} b_1 \right) |z|^2$$

for all $z \in \mathbb{D}$, where $\eta = \min\{\sigma_2, \Gamma_2\}$ and $b_1 = f_{\bar{z}}(0)$.

Proof. Writing $f = h + \bar{g}$ where h and g are given by (1.2), we have

$$\begin{aligned} |f(z)| &\leq (1 + b_1)|z| + \sum_{n=2}^{\infty} (a_n + b_n)|z|^n \\ &\leq (1 + b_1)|z| + \frac{1 - \alpha}{\eta} \sum_{n=2}^{\infty} \left(\frac{\eta}{1 - \alpha} a_n + \frac{\eta}{1 - \alpha} b_n \right) |z|^2 \\ &\leq (1 + b_1)|z| + \frac{1 - \alpha}{\eta} \sum_{n=2}^{\infty} \left(\frac{p_n - \alpha u_n}{1 - \alpha} a_n + \frac{q_n - (-1)^{j-i} \alpha v_n}{1 - \alpha} b_n \right) |z|^2 \\ &\leq (1 + b_1)|z| + \frac{1 - \alpha}{\eta} \left(1 - \frac{q_1 - (-1)^{j-i} \alpha v_1}{1 - \alpha} b_1 \right) |z|^2 \end{aligned}$$

using the hypothesis and applying Theorem 2.4.

The proof of the left hand inequality follows on lines similar to that of the right hand side inequality. $\square$

The covering result for the class $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ follows from the left hand inequality of Theorem 2.7.

Corollary 2.8. Under the hypothesis of Theorem 2.7, we have

$$\left\{ w \in \mathbb{C} : |w| < \frac{1}{\eta} (\eta - 1 + \alpha + (q_1 - (-1)^{j-i} \alpha v_1 - \eta) b_1) \right\} \subset f(\mathbb{D}).$$

Using Theorem 2.4 it is easily seen that the class $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ is convex and closed with respect to the topology of locally uniform convergence so that the closed convex hull of $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ equals itself. The next theorem determines the extreme points of $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$.

Theorem 2.9. *Suppose that $0 \leq \alpha < 1$, $i, j \in \{0, 1\}$, $p_n \geq u_n \geq 0$ ($n = 2, 3, \dots$) and $q_n \geq v_n \geq 0$ ($n = 1, 2, \dots$). Set*

$$h_1(z) = z, \quad h_n(z) = z - \frac{1 - \alpha}{p_n - \alpha u_n} z^n \quad (n = 2, 3, \dots) \quad \text{and}$$

$$g_n(z) = z + \frac{1 - \alpha}{q_n - (-1)^{j-i} \alpha v_n} \bar{z}^n \quad (n = 1, 2, \dots).$$

Then $f \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ if and only if it can be expressed in the form

$$(2.5) \quad f(z) = \sum_{n=1}^{\infty} (X_n h_n + Y_n g_n)(z), \quad X_n \geq 0, \quad Y_n \geq 0 \quad \text{and} \quad \sum_{n=1}^{\infty} (X_n + Y_n) = 1.$$

In particular, the extreme points of $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ are $\{h_n\}$ and $\{g_n\}$.

Proof. Let $f = h + \bar{g} \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ where h and g are given by (1.2). Setting

$$X_n = \frac{p_n - \alpha u_n}{1 - \alpha} a_n \quad (n = 2, 3, \dots) \quad \text{and} \quad Y_n = \frac{q_n - (-1)^{j-i} \alpha v_n}{1 - \alpha} b_n \quad (n = 1, 2, \dots),$$

we note that $0 \leq X_n \leq 1$ ($n = 2, 3, \dots$) and $0 \leq Y_n \leq 1$ ($n = 1, 2, \dots$) by Corollary 2.5. We define

$$X_1 = 1 - \sum_{n=2}^{\infty} X_n - \sum_{n=1}^{\infty} Y_n.$$

By Theorem 2.4, $X_1 \geq 0$ and f can be expressed in the form (2.5).

Conversely, for functions of the form (2.5) we obtain

$$f(z) = z - \sum_{n=2}^{\infty} \mu_n z^n + \sum_{n=1}^{\infty} \nu_n \bar{z}^n,$$

where $\mu_n = (1 - \alpha)X_n/(p_n - \alpha u_n)$ ($n = 2, 3, \dots$) and $\nu_n = (1 - \alpha)Y_n/(q_n - (-1)^{j-i} \alpha v_n)$ ($n = 1, 2, \dots$). Since

$$\sum_{n=2}^{\infty} \frac{p_n - \alpha u_n}{1 - \alpha} \mu_n + \sum_{n=1}^{\infty} \frac{q_n - (-1)^{j-i} \alpha v_n}{1 - \alpha} \nu_n = \sum_{n=2}^{\infty} X_n + \sum_{n=1}^{\infty} Y_n = 1 - X_1 \leq 1,$$

it follows that $f \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ by Theorem 2.4. □

For harmonic functions of the form

$$(2.6) \quad f(z) = z - \sum_{n=2}^{\infty} a_n z^n + \sum_{n=1}^{\infty} b_n \bar{z}^n \quad \text{and} \quad F(z) = z - \sum_{n=2}^{\infty} A_n z^n + \sum_{n=1}^{\infty} B_n \bar{z}^n,$$

where $a_n, b_n, A_n, B_n \geq 0$, we define the product $\hat{*}$ of f and F as

$$(f \hat{*} F)(z) = z - \sum_{n=2}^{\infty} a_n A_n z^n + \sum_{n=1}^{\infty} b_n B_n \bar{z}^n = (F \hat{*} f)(z), \quad z \in \mathbb{D}.$$

Suppose that $\mathcal{I}$ and $\mathcal{J}$ are subclasses of $\mathcal{TH}$. We say that a class $\mathcal{I}$ is closed under $\hat{*}$ if $f\hat{*}F \in \mathcal{I}$ for all $f, F \in \mathcal{I}$. Similarly, the class $\mathcal{I}$ is closed under $\hat{*}$ with members of $\mathcal{J}$ if $f\hat{*}F \in \mathcal{I}$ for all $f \in \mathcal{I}$ and $F \in \mathcal{J}$. In general, the class $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ is not closed under the product $\hat{*}$. The analytic function $f(z) = z - 2z^2$ ($z \in \mathbb{D}$) belongs to $\mathcal{TH}(z + z^2/2, z; 0)$, but $(f\hat{*}f)(z) = z - 4z^2 \notin \mathcal{TH}(z + z^2/2, z; 0)$. However, we shall show that the class $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ is closed under $\hat{*}$ with certain members of $\mathcal{TH}$.

Theorem 2.10. *Suppose that $f, F \in \mathcal{TH}$ are given by (2.6) with $A_n \leq 1$ and $B_n \leq 1$. If $f \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ then $f\hat{*}F \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$.*

Proof. In view of Theorem 2.4, it suffices to show that the coefficients of $f\hat{*}F$ satisfy condition (2.4). Since

$$\begin{aligned} \sum_{n=2}^{\infty} \frac{p_n - \alpha u_n}{1 - \alpha} a_n A_n + \sum_{n=1}^{\infty} \frac{q_n - (-1)^{j-i} \alpha v_n}{1 - \alpha} b_n B_n \\ \leq \sum_{n=2}^{\infty} \frac{p_n - \alpha u_n}{1 - \alpha} a_n + \sum_{n=1}^{\infty} \frac{q_n - (-1)^{j-i} \alpha v_n}{1 - \alpha} b_n \leq 1, \end{aligned}$$

the result follows immediately. $\square$

By imposing some restrictions on the coefficients of Φ_i , it is possible for the class $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ to be closed under the product $\hat{*}$, as seen by the following corollary.

Corollary 2.11. *If $f, F \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ with $p_n \geq 1$ ($n = 2, 3, \dots$) and $q_n \geq 1$ ($n = 1, 2, \dots$) then $f\hat{*}F \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$.*

Proof. Suppose that f and F are given by (2.6). Using the hypothesis and Corollary 2.5 it is easy to see that $A_n \leq 1$ ($n = 2, 3, \dots$) and $B_n \leq 1$ ($n = 1, 2, \dots$). The result now follows by Theorem 2.10. $\square$

For $c > -1$ and a harmonic function $f = h + \bar{g}$, let $L_c[f]$ be the integral operator defined by

$$L_c[f](z) = \frac{c+1}{z^c} \int_0^z t^{c-1} h(t) dt + \overline{\frac{c+1}{z^c} \int_0^z t^{c-1} g(t) dt}, \quad z \in \mathbb{D}.$$

From the representation of $L_c[f]$, it is easy to deduce that

$$L_c[f](z) = f(z) \hat{*} \left(z - \sum_{n=2}^{\infty} \frac{c+1}{c+n} z^n + \sum_{n=1}^{\infty} \frac{c+1}{c+n} \bar{z}^n \right), \quad z \in \mathbb{D}$$

so that Theorem 2.10 gives

Corollary 2.12. *If $f \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ then $L_c[f] \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$.*

Now, we determine the convex combination properties of the members of the class $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$.

Theorem 2.13. *The class $\mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ is closed under convex combinations.*

Proof. For $k = 1, 2, \dots$ suppose that $f_k \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ where

$$f_k(z) = z - \sum_{n=2}^{\infty} a_{n,k} z^n + \sum_{n=1}^{\infty} b_{n,k} \bar{z}^n, \quad z \in \mathbb{D}.$$

For $\sum_{k=1}^{\infty} t_k = 1$, $0 \leq t_k \leq 1$, the convex combination of f_k 's may be written as

$$f(z) = \sum_{k=1}^{\infty} t_k f_k(z) = z - \sum_{n=2}^{\infty} \gamma_n z^n + \sum_{n=1}^{\infty} \delta_n \bar{z}^n, \quad z \in \mathbb{D}$$

where $\gamma_n = \sum_{k=1}^{\infty} t_k a_{n,k}$ ($n = 2, 3, \dots$) and $\delta_n = \sum_{k=1}^{\infty} t_k b_{n,k}$ ($n = 1, 2, \dots$). Since

$$\begin{aligned} & \sum_{n=2}^{\infty} \frac{p_n - \alpha u_n}{1 - \alpha} \gamma_n + \sum_{n=1}^{\infty} \frac{q_n - (-1)^{j-i} \alpha v_n}{1 - \alpha} \delta_n \\ &= \sum_{k=1}^{\infty} t_k \left(\sum_{n=2}^{\infty} \frac{p_n - \alpha u_n}{1 - \alpha} a_{n,k} + \sum_{n=1}^{\infty} \frac{q_n - (-1)^{j-i} \alpha v_n}{1 - \alpha} b_{n,k} \right) \\ &\leq \sum_{k=1}^{\infty} t_k = 1, \end{aligned}$$

it follows that $f \in \mathcal{TH}(\Phi_i, \Psi_j; \alpha)$ by Theorem 2.4. $\square$

3. AN APPLICATION IN UNIVALENT HARMONIC FUNCTION THEORY

For $0 \leq \alpha < 1$, let $\mathcal{U}_H(\alpha)$ denote the set of all harmonic functions $f \in \mathcal{H}$ that satisfy $\operatorname{Re} f(z)/z > \alpha$ for $z \in \mathbb{D}$ and let $\mathcal{TU}_H(\alpha) := \mathcal{U}_H(\alpha) \cap \mathcal{TH}$. Setting $\Phi(z) = z/(1-z) + \bar{z}/(1-\bar{z})$ in Corollary 2.6 and using the results of Section 2, we obtain

Theorem 3.1. *Let the function $f = h + \bar{g}$ be such that h and g are given by (1.2). Then $f \in \mathcal{TU}_H(\alpha)$ if and only if*

$$\sum_{n=2}^{\infty} \frac{a_n}{1 - \alpha} + \sum_{n=1}^{\infty} \frac{b_n}{1 - \alpha} \leq 1.$$

Furthermore, if $f \in \mathcal{TU}_H(\alpha)$ then $a_n \leq 1 - \alpha$ ($n = 2, 3, \dots$), $b_n \leq 1 - \alpha$ ($n = 1, 2, \dots$) and

$$(3.1) \quad (1 - b_1)|z| - (1 - \alpha - b_1)|z|^2 \leq |f(z)| \leq (1 + b_1)|z| + (1 - \alpha - b_1)|z|^2$$

for all $z \in \mathbb{D}$. In particular, the range $f(\mathbb{D})$ contains the disk $|w| < \alpha$.

Moreover, the extreme points of the class $\mathcal{TU}_H(\alpha)$ are $\{h_n\}$ and $\{g_n\}$ where $h_1(z) = z$, $h_n(z) = z - (1 - \alpha)z^n$ ($n = 2, 3, \dots$) and $g_n(z) = z + (1 - \alpha)\bar{z}^n$ ($n = 1, 2, \dots$).

The upper bound given in (3.1) for $f \in \mathcal{TU}_H(\alpha)$ is sharp and equality occurs for the function $f(z) = z + b_1 \bar{z} + (1 - \alpha - b_1) \bar{z}^2$ for $b_1 \leq 1 - \alpha$. By Corollary 2.11, the class $\mathcal{TU}_H(\alpha)$ is closed under the product $\hat{*}$. In fact, we have

$$\mathcal{TU}_H(\alpha) \hat{*} \mathcal{TU}_H(\beta) \subset \mathcal{TU}_H(1 - (1 - \alpha)(1 - \beta))$$

for $\alpha, \beta \in [0, 1)$.

Making use of Theorem 3.1 and the well-known Gauss's summation theorem: if $\operatorname{Re}(c - a - b) > 0$ then

$$F(a, b, c; 1) = \frac{\Gamma(c)\Gamma(c - a - b)}{\Gamma(c - a)\Gamma(c - b)}, \quad c \neq 0, -1, -2, \dots$$

we have the following corollary.

Corollary 3.2. *Let $a_k, b_k > 0$, $c_k > a_k + b_k + 1$ for $k = 1, 2$. Furthermore, if*

$$F(a_1, b_1, c_1; 1) + F(a_2, b_2, c_2; 1) \leq 3$$

then $\phi_1 + \bar{\phi}_2 \in \mathcal{U}_H(\alpha)$, ϕ_1 and ϕ_2 being given by (2.3).

For further investigation of results regarding $\mathcal{TU}_H(\alpha)$, we need to prove the following simple lemma.

Lemma 3.3. *Let $f = h + \bar{g} \in \mathcal{H}$ where h and g are given by (1.1) with $b_1 = g'(0) = 0$. Suppose that $\lambda \in (0, 1]$.*

- (i) *If $\sum_{n=2}^{\infty} (|a_n| + |b_n|) \leq \lambda$ then $f \in \mathcal{U}_H(1 - \lambda)$;*
- (ii) *If $\sum_{n=2}^{\infty} n(|a_n| + |b_n|) \leq \lambda$ then $f \in \mathcal{U}_H(1 - \lambda/2)$ and is starlike of order $2(1 - \lambda)/(2 + \lambda)$.*

The results are sharp.

Proof. Part (i) follows by Theorem 3.1. For the proof of (ii), note that

$$\sum_{n=2}^{\infty} (|a_n| + |b_n|) \leq \frac{1}{2} \sum_{n=2}^{\infty} n(|a_n| + |b_n|) \leq \frac{\lambda}{2}.$$

By part (i) of the theorem, $f \in \mathcal{U}_H(1 - \lambda/2)$. The order of starlikeness of f follows by [15]. The functions $z + \lambda\bar{z}^2$ and $z + \lambda\bar{z}^2/2$ show that the results in (i) and (ii) respectively are best possible. $\square$

A well-known classical result involving differential inequalities in univalent function theory is Marx Stroh acker theorem which states that if f is an analytic function in $\mathbb{D}$ with $f(0) = 0 = f'(0) - 1$ then

$$\operatorname{Re} \left(1 + \frac{zf''(z)}{f'(z)} \right) > 0 \quad \Rightarrow \quad \operatorname{Re} \frac{zf'(z)}{f(z)} > \frac{1}{2} \quad \Rightarrow \quad \operatorname{Re} \frac{f(z)}{z} > \frac{1}{2} \quad (z \in \mathbb{D}).$$

The function $f(z) = z/(1 - z)$ shows that all these implications are sharp. However, this theorem does not extend to univalent harmonic mappings, that is, if $f \in \mathcal{S}_H$ maps $\mathbb{D}$ onto a convex domain then it is not true in general that $\operatorname{Re} f(z)/z > 0$ for all $z \in \mathbb{D}$. To see this, consider the harmonic half-plane mapping

$$L(z) = \frac{z - \frac{1}{2}z^2}{(1 - z)^2} + \overline{\frac{-\frac{1}{2}z^2}{(1 - z)^2}}, \quad z \in \mathbb{D}$$

which maps the unit disk $\mathbb{D}$ univalently onto the half-plane $\operatorname{Re} w > -1/2$. Figure 1 shows that the function $L(z)/z$ does not have a positive real part in $\mathbb{D}$.

Denote by $\mathcal{TS}_H^{*0}(\alpha)$, $\mathcal{TK}_H^0(\alpha)$ and $\mathcal{TU}_H^0(\alpha)$, the class consisting of functions f in $\mathcal{TS}_H^*(\alpha)$, $\mathcal{TK}_H(\alpha)$ and $\mathcal{TU}_H(\alpha)$ respectively, for which $f_{\bar{z}}(0) = 0$. The next theorem connects the relation between these three classes.

Theorem 3.4. For $0 \leq \alpha < 1$, the following sharp inclusions hold:

$$(3.2) \quad \mathcal{TK}_H^0(\alpha) \subset \mathcal{TU}_H^0\left(\frac{3-\alpha}{2(2-\alpha)}\right);$$

and

$$(3.3) \quad \mathcal{TS}_H^{*0}(\alpha) \subset \mathcal{TU}_H^0\left(\frac{1}{2-\alpha}\right).$$

Proof. Let $f = h + \bar{g} \in \mathcal{TH}$ where h and g are given by (1.2). If $f \in \mathcal{TK}_H^0(\alpha)$ then

$$\sum_{n=2}^{\infty} n(a_n + b_n) \leq \frac{1}{2-\alpha} \sum_{n=2}^{\infty} n(n-\alpha)(a_n + b_n) \leq \frac{1-\alpha}{2-\alpha}$$

using [8]. By Lemma 3.3(ii) $f \in \mathcal{TU}_H^0((3-\alpha)/(2(2-\alpha)))$. Regarding the other inclusion, note that if $f \in \mathcal{TS}_H^{*0}(\alpha)$ then

$$\sum_{n=2}^{\infty} (a_n + b_n) \leq \frac{1}{2-\alpha} \sum_{n=2}^{\infty} (n-\alpha)(a_n + b_n) \leq \frac{1-\alpha}{2-\alpha}$$

by [9]. This shows that $f \in \mathcal{TU}_H^0(1/(2-\alpha))$ by Lemma 3.3(i) as desired. The analytic functions $z - (1-\alpha)z^2/(2(2-\alpha))$ and $z - (1-\alpha)z^2/(2-\alpha)$ show that inclusions in (3.2) and (3.3) respectively are sharp. $\square$

Remark 3.5. The proof of Theorem 3.4 shows that if $f \in \mathcal{TK}_H^0(\alpha)$ then f is starlike of order $2/(5-3\alpha)$ by applying Lemma 3.3(ii). This gives the inclusion

$$\mathcal{TK}_H^0(\alpha) \subset \mathcal{TS}_H^{*0}\left(\frac{2}{5-3\alpha}\right).$$

It is not known whether this inclusion is sharp for $\alpha \in (0, 1)$. However, if $\alpha = 0$ then the inclusion $\mathcal{TK}_H^0(0) \subset \mathcal{TS}_H^{*0}(2/5)$ is sharp with the extremal function as $f(z) = z + \bar{z}^2/4$.




FIGURE 1. Graph of the function $L(z)/z$.

The functions in the class $\mathcal{TU}_H(\alpha)$ need not be univalent in $\mathbb{D}$. The last theorem of this section determines the radius of univalence, starlikeness and convexity of the class $\mathcal{TU}_H^0(\alpha)$.

Theorem 3.6. *The radius of univalence of the class $\mathcal{TU}_H^0(\alpha)$ is $1/(2(1-\alpha))$. This bound is also the radius of starlikeness of $\mathcal{TU}_H^0(\alpha)$. The radius of convexity of the class $\mathcal{TU}_H^0(\alpha)$ is $1/(4(1-\alpha))$.*

Proof. Let $f = h + \bar{g} \in \mathcal{TU}_H^0(\alpha)$ where h and g are given by (1.2) and let $r \in (0, 1)$ be fixed. Then $r^{-1}f(rz) \in \mathcal{TU}_H^0(\alpha)$ and we have

$$\sum_{n=2}^{\infty} n(a_n + b_n)r^{n-1} \leq \sum_{n=2}^{\infty} \left(\frac{a_n}{1-\alpha} + \frac{b_n}{1-\alpha} \right) \leq 1$$

provided $nr^{n-1} \leq 1/(1-\alpha)$ which is true if $r \leq 1/(2(1-\alpha))$. In view of [17, Theorem 1, p. 284], f is univalent and starlike in $|z| < 1/(2(1-\alpha))$. Regarding the radius of convexity, note that

$$\sum_{n=2}^{\infty} n^2(a_n + b_n)r^{n-1} \leq \sum_{n=2}^{\infty} \left(\frac{a_n}{1-\alpha} + \frac{b_n}{1-\alpha} \right) \leq 1$$

provided $n^2r^{n-1} \leq 1/(1-\alpha)$ which is true if $r \leq 1/(4(1-\alpha))$. The function $f(z) = z + (1-\alpha)\bar{z}^2$ shows that these bounds are sharp. This completes the proof of the theorem. $\square$

REFERENCES

- [1] O. P. Ahuja and J. M. Jahangiri, Noshiro-type harmonic univalent functions, *Sci. Math. Japon.*, **56** (2) (2002), 293–299.
- [2] O. P. Ahuja and J. M. Jahangiri, A subclass of harmonic univalent functions, *J. Nat. Geom.*, **20** (2001), 45–56.
- [3] O. P. Ahuja and H. Silverman, Inequalities associating hypergeometric functions with planer harmonic mappings, *J. Ineq. Pure Appl. Math.* **5** (4) (2004), 1–10.
- [4] J. Clunie and T. Sheil-Small, Harmonic univalent functions, *Ann. Acad. Sci. Fenn. Ser. A. I* **9** (1984), 3–25.
- [5] K. K. Dixit, A. L. Pathak, S. Porwal and R. Agarwal, On a subclass of harmonic univalent functions defined by convolution and integral convolution, *Int. J. Pure Appl. Math.*, **69** (3) (2011), 255–264.
- [6] K. K. Dixit and S. Porwal, Some properties of harmonic functions defined by convolution, *Kyungpook Math. J.*, **49** (2009), 751–761.
- [7] B. A. Frasin, Comprehensive family of harmonic univalent functions, *SUT J. Math.*, **42** (1) (2006), 145–155.
- [8] J. M. Jahangiri, Coefficient bounds and univalence criteria for harmonic functions with negative coefficients, *Ann. Univ. Mariae Cruie-Sklodowska Sect. A* **52** (1998), no.2, 57–66.
- [9] J. M. Jahangiri, Harmonic functions starlike in the unit disk, *J. Math. Anal. Appl.* **235** (1999), 470–477.
- [10] S. S. Joshi, Subclasses of harmonic univalent functions associated with hypergeometric functions, *Int. J. Pure Appl. Math.*, **60** (1) (2010), 5–14.
- [11] O. P. Juneja, T. R. Reddy and M. L. Mogra, A convolution approach for analytic functions with negative coefficients, *Soochow J. Math.*, **11** (1985), 69–81.
- [12] G. Murugusundaramoorthy and R. K. Raina, On a subclass of harmonic functions associated with the Wright's generalized hypergeometric functions, *Hacet. J. Math. Stat.* **38** (2) (2009), 129–136.
- [13] G. Murugusundaramoorthy, K. Uma and M. Acu, Harmonic functions prestarlike in the unit disc, *Gen. Math.*, **18** (4) (2010), 149–161.

- [14] G. Murugusundaramoorthy and K. Uma, Harmonic univalent functions associated with generalized hypergeometric functions, *Bull. Math. Anal. Appl.*, **2** (2) (2010), 69–76.
- [15] S. Nagpal and V. Ravichandran, A subclass of close-to-convex harmonic mappings, *Complex Var. Elliptic Equ.* (2012), DOI: 10.1080/17476933.2012.727409.
- [16] S. Nagpal and V. Ravichandran, Fully starlike and fully convex harmonic mappings of order α , *Annales Polonici Mathematici* (2012), To appear.
- [17] H. Silverman, Harmonic univalent functions with negative coefficients, *J. Math. Anal. Appl.* **220**(1998), 283–289.
- [18] H. Silverman and E. M. Silvia, Subclasses of harmonic univalent functions, *New Zealand J. Math.*, **28** (1999) 275–284.
- [19] S. Yalçın and M. Öztürk, On a subclass of certain convex harmonic functions, *J. Korean Math. Soc.*, **43** (4) (2006), 803–813.
- [20] S. Yalçın, M. Öztürk and M. Yamankaradeniz, Convex subclass of harmonic starlike functions, *Appl. Math. Comput.*, **154** (2004), 449–459.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DELHI, DELHI–110 007, INDIA
E-mail address: sumitnagpal.du@gmail.com

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DELHI, DELHI–110 007, INDIA
E-mail address: vravi68@gmail.com